

# Latency Period of Malignant Lymphoma in Mice Exposed to Continuous Low-dose-rate Gamma-rays – Pathological Study –

Satoshi TANAKA\*, Ignacia TANAKA\*, Keiji OGURA\*, Kazuaki ICHINOHE\*,  
Yoichi OGHISO\*, Fumiaki SATO\*\*, Kimio TANAKA\*  
\*Department of Radiobiology, \*\*General Advisor

## Abstract

Late effects of low-dose and low-dose-rate (LDR) ionizing radiation are potential hazards, and they have become a serious concern since the nuclear accident at the Fukushima Dai-ichi Nuclear Power Plant. Chronic exposure of 4000 mice to LDR gamma rays showed that the average life span of female mice irradiated at 21 mGy/22 h/day (909  $\mu$ Gy/h) for 400 days (total dose: 8000 mGy) were shortened by about 120 days due to premature death from various neoplasms including malignant lymphomas. This suggest that long-term exposure to LDR gamma-rays caused early onset or increase progression of neoplasms. A cross-sectional study to observe the development and progression of neoplasia and non-neoplastic lesions is underway.

Preliminary findings showed increased incidences ( $p<0.05$ ) and early emergence of malignant tumors of liver and lung origins in the irradiated group (20 mGy/22 h/day). Malignant lymphomas and benign tumors of the liver appeared at the same age in both irradiated and non-irradiated groups. The numbers of non-neoplastic lesions (ovarian atrophy) increased significantly ( $p<0.01$ ) in mice exposed for 200 days (total dose: 4000 mGy). These suggested that the effects of LDR irradiation on neoplastic and non-neoplastic lesions varied depending on the organ and/or tissue.




Fig. 1 Incidence of (a) Liver, Hepatocellular adenoma, (b) Liver, Hepatocellular carcinoma, (c) Lung, Bronchiolo-Alveolar adenoma, (d) Lung, Bronchiolo-Alveolar carcinoma, (e) Lymphoma, Malignant and (f) Ovarian atrophy (\*: $P<0.05$ , \*\*:  $P<0.01$ ).